

FORM PTO-4390 (Modified)
(REV 11-2000)

U.S. DEP.

COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES

DESIGNATED/ELECTED OFFICE (DO/EO/US)

CONCERNING A FILING UNDER 35 U.S.C. 371

PA-9848

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

To be assigned 09/869629

INTERNATIONAL APPLICATION NO.

PCT/GB99/04410

INTERNATIONAL FILING DATE

December 23, 1999

PRIORITY DATE CLAIMED

December 30, 1998

TITLE OF INVENTION

NMR Spectroscopic In Vitro Assay Using Hyperpolarization

APPLICANT(S) FOR DO/EO/US

P. Knox, et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☒ is attached hereto (required only if not communicated by the International-Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☒ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

copy of this transmittal letter for charging purposes

copy of the International Application as published by the International Bureau

return postcard

APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.492(a)(1) - (5)) :

09/069629

NATIONAL APPLICATION NO.

PCT/GB99/04410

ATTORNEY'S DOCKET NUMBER

PA-9848

The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :

- ☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000.00
- ☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$860.00

Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)).

☐ 20

☐ 30

\$0.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	29 - 20 =	9	x \$18.00
Independent claims	1 - 3 =	0	x \$80.00

\$162.00

\$0.00

Multiple Dependent Claims (check if applicable).

☐

\$0.00

TOTAL OF ABOVE CALCULATIONS =

\$1,022.00

☐ Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2.

\$0.00

SUBTOTAL =

\$1,022.00

Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)).

☐ 20

☐ 30

\$0.00

TOTAL NATIONAL FEE =

\$1,022.00

Fees for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).

☐

\$0.00

TOTAL FEES ENCLOSED =

\$1,022.00

Amount to be:

refunded

\$

charged

\$


- ☐ A check in the amount of _____ to cover the above fees is enclosed.
- ☒ Please charge my Deposit Account No. 500-588 in the amount of \$1,022.00 to cover the above fees. A duplicate copy of this sheet is enclosed.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 500-588. A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Royal N. Ronning, Jr.
Amersham Pharmacia Biotech, Inc.
800 Centennial Avenue
Piscataway, New Jersey 08855

(732) 457-8423


SIGNATURE

Royal N. Ronning, Jr.

NAME

32,529

REGISTRATION NUMBER

June 28, 2001

DATE

PA-9848

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: P. Knox, et al. Group Art Unit: To be assigned
Serial Number: To be assigned Examiner: To be assigned
Filing Date: June 28, 2001
Title: NMR Spectroscopic In Vitro Assay Using Hyperpolarization

FIRST PRELIMINARY AMENDMENT

Honorable Assistant Commissioner of Patents
Box New Patent Application
Washington, D.C. 20231

Sir:

Please consider the following amendments and remarks in connection with the prosecution of the captioned application, which is a filing under 35 U.S.C. § 371 and claims priority to international application number PCT/GB99/04410 filed December 23, 1999, application number 9828852.5 filed in Great Britain on December 30, 1998, and application number 9918096.0 filed in Great Britain on August 2, 1999.

In the Claims

Please amend page 21, line 1 as follows:

[C L A I M S]

What is claimed is:

Please amend claim 4 as follows:

4. (once amended) The method of [any of claims 1 to 3]claim 1, wherein the assay reagent is a compound which contains an artificially high concentration of an NMR active nucleus.

Please amend claim 6 as follows:

6. (once amended) The method of [any of claims 1 to 5]claim 1, wherein the assay reagent is an organic compound comprising one or more NMR active nuclei associated with a bond which is broken during the course of the assay.

Please amend claim 8 as follows:

8. (once amended) The method of [any claim 1-7]claim 1, wherein the assay reagent is analysed repeatedly in step c) at known time intervals so as to generate information about a change with time of the assay reagent.

Please amend claim 9 as follows:

9. (once amended) The method of [any one claim 1 to 8]claim 1, wherein the [asseay]assay reagent is a Nucleotide, or nucleotide analogue, polynucleotide, amino acid analogue, polypeptide or protein.

Please amend claim 10 as follows:

10. (once amended) The method of [any one of claims 1 to 9]claim 1, wherein the assay is a nucleic acid hybridisation assay.

Please amend claim 11 as follows:

11. (once amended) The method of [any one of claims 1 to 10]claim 1, wherein the assay is a binding assay.

Please amend claim 12 as follows:

12. (once amended) The method of [claims 1 to 11]claim 1, wherein the assay reagent is a compound specifically labelled with at least one NMR active nucleus and the assay reagent is administered to a micro-organism, macro-organism or cultured cells, cellular metabolites or an excretion product of the assay reagent are hyperpolarised and analysed by nuclear magnetic resonance spectroscopy, nuclear magnetic resonance imaging or both.

Please amend claim 13 as follows:

13. (once amended) The method of [claims 1 to 12]claim 1, wherein the assay is a binding study performed using micro-organisms or cultured cells

Please amend claim 14 as follows:

14. (once amended) The method of [claims 1 to 13]claim 1 wherein the hyperpolarisation transfer is repeated to enhance the signal-to-noise ratio.

Please amend claim 15 as follows:

15. (once amended) The method of [claim 1 to 14]claim 1 wherein the shortening

effect as expressed by the improvement of signal-to-noise per unit time is a factor of 10 or more compared to known assay techniques without hyperpolarisation.

Please amend claim 16 as follows:

16. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer from a hyperpolarised noble gas, or a mixture of hyperpolarised noble gases.

Please amend claim 19 as follows:

19. (once amended) The method of [claims 16 to 18]claim 16 wherein the hyperpolarisation is transferred by a hyperpolarised noble gas in solution and wherein the viscosity of the solution is at least 1000 mPs.

Please amend claim 20 as follows:

20. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer at a temperature of 4.2 K or less in the presence of a magnetic field of at least 1 T.

Please amend claim 21 as follows:

21. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer using dynamic nuclear polarisation.

Please amend claim 22 as follows:

22. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by para hydrogen induced polarisation.

Please amend claim 23 as follows:

23. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out with the spin refrigeration technique.

Please amend claim 24 as follows:

24. (once amended) The method of [claims 1 to 23]claim 1, wherein more than one

assay is multiplexed and monitored by NMR spectroscopy and/or NMR imaging.

Please amend claim 25 as follows:

25. (once amended) The method of [claims 1 to 24]claim 1 wherein the assay is performed in a multiwell or multispot assay array.

Please amend claim 26 as follows:

26. (once amended) The method of [claims 1 to 25]claim 1 wherein step c) is performed by examining the assay reagent using both NMR spectroscopy to obtain more than one spectrum, and magnetic resonance imaging to obtain one or more discrete spectral location, and repeating the examination at least once so as to obtain quantitative information about kinetic or time-dependant alteration in chemistry, environment or structure of the assay reagent.

Please amend claim 27 as follows:

27. (once amended) The method of [claim 1 to 26]claim 1, wherein step c) is performed in an aerosol or flow-through device applied to aerosol droplets where the well, surface or container is used to contain the assay reagent.

Please amend claim 28 as follows:

28. (once amended) An *in vitro* assay kit for carrying out the assay method as defined in claim 1 [to 27]which comprises: one or more assay reagents each containing at least one NMR active nucleus contained in a well or vial or other suitable container for carrying out the hyperpolarisation of step (b) of claim 1.

Please amend claim 29 as follows:

29. (once amended) The *in vitro* kit of claim 28 where the NMR analysis of step (c) [of claim 1] is carried out in the same well, vial or container as the hyperpolarisation transfer is carried out.

Remarks

Claims 1-29 are pending in the instant application. Applicants have amended claims 4, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, and 29 to more fully conform with U.S. practice and to delete multiple dependencies. A version of the claims marked up to show the amendments, as well as a clean version of the claims encompassing the amendments, is attached hereto.

Applicants respectfully assert that all amendments are fairly based on the specification, and respectfully request their entry.

Applicants believe that the claims, as amended, are in allowable form, and earnestly solicit the allowance of claims 1-29.

Respectfully submitted,

A handwritten signature in black ink, appearing to read "Royal N. Ronning, Jr.", written in a cursive style.

Royal N. Ronning, Jr. 32,529
Attorney for Applicants

Amersham Pharmacia Biotech, Inc.
800 Centennial Avenue
P. O. Box 1327
Piscataway, New Jersey 08855-1327

Tel: (732) 457-8423
Fax: (732) 457-8463

Claims (marked-up version showing amendment(s))

[C L A I M S]

What is claimed is:

4. (once amended) The method of [any of claims 1 to 3]claim 1, wherein the assay reagent is a compound which contains an artificially high concentration of an NMR active nucleus.
6. (once amended) The method of [any of claims 1 to 5]claim 1, wherein the assay reagent is an organic compound comprising one or more NMR active nuclei associated with a bond which is broken during the course of the assay.
8. (once amended) The method of [any claim 1-7]claim 1, wherein the assay reagent is analysed repeatedly in step c) at known time intervals so as to generate information about a change with time of the assay reagent.
9. (once amended) The method of [any one claim 1 to 8]claim 1, wherein the [asseay]assay reagent is a Nucleotide, or nucleotide analogue, polynucleotide, amino acid analogue, polypeptide or protein.
10. (once amended) The method of [any one of claims 1 to 9]claim 1, wherein the

assay is a nucleic acid hybridisation assay.

11. (once amended) The method of [any one of claims 1 to 10]claim 1, wherein the assay is a binding assay.
12. (once amended) The method of [claims 1 to 11]claim 1, wherein the assay reagent is a compound specifically labelled with at least one NMR active nucleus and the assay reagent is administered to a micro-organism, macro-organism or cultured cells, cellular metabolites or an excretion product of the assay reagent are hyperpolarised and analysed by nuclear magnetic resonance spectroscopy, nuclear magnetic resonance imaging or both.
13. (once amended) The method of [claims 1 to 12]claim 1, wherein the assay is a binding study performed using micro-organisms or cultured cells
14. (once amended) The method of [claims 1 to 13]claim 1 wherein the hyperpolarisation transfer is repeated to enhance the signal-to-noise ratio.
15. (once amended) The method of [claim 1 to 14]claim 1 wherein the shortening effect as expressed by the improvement of signal-to-noise per unit time is a factor of 10 or more compared to known assay techniques without hyperpolarisation.

16. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer from a hyperpolarised noble gas, or a mixture of hyperpolarised noble gases.
19. (once amended) The method of [claims 16 to 18]claim 16 wherein the hyperpolarisation is transferred by a hyperpolarised noble gas in solution and wherein the viscosity of the solution is at least 1000 mPs.
20. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer at a temperature of 4.2 K or less in the presence of a magnetic field of at least 1 T.
21. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer using dynamic nuclear polarisation.
22. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by para hydrogen induced polarisation.

23. (once amended) The method of [claims 1 to 15]claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out with the spin refrigeration technique.
24. (once amended) The method of [claims 1 to 23]claim 1, wherein more than one assay is multiplexed and monitored by NMR spectroscopy and/or NMR imaging.
25. (once amended) The method of [claims 1 to 24]claim 1 wherein the assay is performed in a multiwell or multispot assay array.
26. (once amended) The method of [claims 1 to 25]claim 1 wherein step c) is performed by examining the assay reagent using both NMR spectroscopy to obtain more than one spectrum, and magnetic resonance imaging to obtain one or more discrete spectral location, and repeating the examination at least once so as to obtain quantitative information about kinetic or time-dependant alteration in chemistry, environment or structure of the assay reagent.
27. (once amended) The method of [claim 1 to 26]claim 1, wherein step c) is performed in an aerosol or flow-through device applied to aerosol droplets where the well, surface or container is used to contain the assay reagent.
28. (once amended) An *in vitro* assay kit for carrying out the assay method as defined

in claim 1 [to 27]which comprises: one or more assay reagents each containing at least one NMR active nucleus contained in a well or vial or other suitable container for carrying out the hyperpolarisation of step (b) of claim 1.

29. (once amended) The *in vitro* kit of claim 28 where the NMR analysis of step (c) [of claim 1] is carried out in the same well, vial or container as the hyperpolarisation transfer is carried out.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219

Claims (clean version encompassing amendments)

What is claimed is:

1. An *in vitro* assay method which comprises:
 - a) using an assay reagent containing at least one NMR active nucleus to perform an assay, and
 - b) hyperpolarising at least one NMR active nucleus of the assay reagent;
wherein steps (a) and (b) are performed simultaneously or sequentially in either order, and
 - c) analysing the assay reagent and/or the assay by NMR and
 - d) optionally using the NMR data obtained in step c) to generate further assay result(s).
2. The method of claim 1 wherein the NMR active nucleus ^{15}N , ^{19}F , ^{31}P , ^1H , ^{29}Si and/or ^{13}C .
3. The method of claim 2, wherein the NMR active nucleus is ^{15}N or ^{13}C .
4. (once amended) The method of claim 1, wherein the assay reagent is a compound which contains an artificially high concentration of an NMR active nucleus.

5. The method of claim 4, wherein the assay reagent contains an artificially high concentration in 1-10 defined positions.
6. (once amended) The method of claim 1, wherein the assay reagent is an organic compound comprising one or more NMR active nuclei associated with a bond which is broken during the course of the assay.
7. The method of claim 6, wherein the assay reagent contains two or more NMR active nuclei and each NMR active nucleus produces a distinct NMR spectrum and when the assay method is performed, it results in changes to the chemical and/or physical environment of the nucleus and this is mirrored by spectral changes which can be monitored.
8. (once amended) The method of claim 1, wherein the assay reagent is analysed repeatedly in step c) at known time intervals so as to generate information about a change with time of the assay reagent.
9. (once amended) The method of claim 1, wherein the assay reagent is a Nucleotide, or nucleotide analogue, polynucleotide, amino acid analogue, polypeptide or protein.
10. (once amended) The method of claim 1, wherein the assay is a nucleic acid

hybridisation assay.

11. (once amended) The method of claim 1, wherein the assay is a binding assay.
12. (once amended) The method of claim 1, wherein the assay reagent is a compound specifically labelled with at least one NMR active nucleus and the assay reagent is administered to a micro-organism, macro-organism or cultured cells, cellular metabolites or an excretion product of the assay reagent are hyperpolarised and analysed by nuclear magnetic resonance spectroscopy, nuclear magnetic resonance imaging or both.
13. (once amended) The method of claim 1, wherein the assay is a binding study performed using micro-organisms or cultured cells
14. (once amended) The method of claim 1 wherein the hyperpolarisation transfer is repeated to enhance the signal-to-noise ratio.
15. (once amended) The method of claim 1 wherein the shortening effect as expressed by the improvement of signal-to-noise per unit time is a factor of 10 or more compared to known assay techniques without hyperpolarisation.
16. (once amended) The method of claim 1 where the hyperpolarisation of the NMR

active nucleus of the assay reagent is carried out by polarisation transfer from a hyperpolarised noble gas, or a mixture of hyperpolarised noble gases.

17. The method of claim 16 wherein the noble gas is ^{129}Xe .
18. The method of claim 16 wherein the noble gas is ^3He .
19. (once amended) The method of claim 16 wherein the hyperpolarisation is transferred by a hyperpolarised noble gas in solution and wherein the viscosity of the solution is at least 1000 mPs.
20. (once amended) The method of claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer at a temperature of 4.2 K or less in the presence of a magnetic field of at least 1 T.
21. (once amended) The method of claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer using dynamic nuclear polarisation.
22. (once amended) The method of claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by para hydrogen induced polarisation.

23. (once amended) The method of claim 1 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out with the spin refrigeration technique.
24. (once amended) The method of claim 1, wherein more than one assay is multiplexed and monitored by NMR spectroscopy and/or NMR imaging.
25. (once amended) The method of claim 1 wherein the assay is performed in a multiwell or multispot assay array.
26. (once amended) The method of claim 1 wherein step c) is performed by examining the assay reagent using both NMR spectroscopy to obtain more than one spectrum, and magnetic resonance imaging to obtain one or more discrete spectral location, and repeating the examination at least once so as to obtain quantitative information about kinetic or time-dependant alteration in chemistry, environment or structure of the assay reagent.
27. (once amended) The method of claim 1, wherein step c) is performed in an aerosol or flow-through device applied to aerosol droplets where the well, surface or container is used to contain the assay reagent.

28. (once amended) An *in vitro* assay kit for carrying out the assay method as defined in claim 1 which comprises: one or more assay reagents each containing at least one NMR active nucleus contained in a well or vial or other suitable container for carrying out the hyperpolarisation of step (b) of claim 1.
29. (once amended) The *in vitro* kit of claim 28 where the NMR analysis of step (c) is carried out in the same well, vial or container as the hyperpolarisation transfer is carried out.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221

NMR SPECTROSCOPIC IN VITRO ASSAY USING HYPERPOLARIZATION

TECHNICAL FIELD

5 The present invention relates to tailoring the shape of a magneto-resistive material, and more particularly to a design of the shape of the magneto-resistive material to obtain a new type of position sensitive sensor.

BACKGROUND

10 The position of a moving object is often determined by means of the readout from a resistive sensor, usually of potentiometer type, which is mechanically connected to the object to be monitored.

15 In order to reduce the wear and thereby increase the reliability, it is desirable to eliminate the sliding friction encountered in the standard resistive sensors. Non-contact methods using e.g. inductively coupled coils is currently being introduced as replacement for the potentiometer sensors. However, these are more complex and therefore more expensive.

20 In recent years novel types of magneto-resistive materials with much higher sensitivity to moderate changes in magnetic fields have been found. These new materials showing giant magneto-resistance (GMR) or colossal magneto-resistance (CMR) make possible new types of position sensors.

25 In a document U.S. Patent No. 5,475,304 is disclosed a giant magneto-resistant sensor including at least one layered structure. The layered structure includes a ferromagnetic layer having a fixed magnetic state, a second, softer magnetic layer, and a metal layer interposed between and contacting these two layers. The sensor also includes one or more indexing magnets for inducing a domain wall, at a measured position, between regions of nonaligned magnetic fields in the softer magnetic layer. By
30 measuring the resistance across the magneto-resistant sensor a displacement of one workpiece, carrying the sensor, will be measured relative to another workpiece carrying an inducing means.

nuclear polarisation of an assay reagent comprising at least one NMR active nucleus other than the noble gas. The hyperpolarisation of the assay reagent may also be achieved by using an artificially enriched hyperpolarised noble gas, preferably ^3He or ^{129}Xe .

5 Alternatively, hyperpolarisation may be imparted to atoms of significance in biological systems (e.g. ^{13}C , ^{15}N , ^{31}P , ^{29}Si , ^{19}F and ^1H isotopes) by thermodynamic equilibration at very low temperature, suitably below 1K, preferably as close to 0 K as possible, and in the presence of a high magnetic field ("Brute force").

10 A further alternative is that hyperpolarisation may be imparted by dynamic nuclear polarisation (DNP). In the solid phase, the material is mixed with a paramagnetic species (DNP agent), for example a transition metal ion such as chromium (V) or manganese (II) and/or a free radical generator or other particles having associated free electrons. The method utilises a moderate or high magnetic field and very low temperature, e.g. by carrying out the conversion in
15 liquid helium and a magnetic field of about 1 T or above.

A further technique for imparting hyperpolarisation is para hydrogen induced polarisation which involves cooling hydrogen to a low temperature, e.g. 20 K or less, to give para hydrogen enriched hydrogen. This enriched hydrogen is then used to hydrogenate an unsaturated
20 target organic molecule (containing NMR active nuclei) imparting a non-thermodynamic spin configuration to the target molecule.

A yet further method covered by the present invention for preparing hyperpolarised materials is spin refrigeration. With this technique, the assay reagent is doped with or intimately
25 mixed with a suitable paramagnetic material in crystal form (e.g. crystalline powder) with a symmetry axis of order three or more. One advantage with this technique is that there is no need for a uniform magnetic field since no resonant excitation field is applied. The sample is rotated to bring the electron paramagnetic resonance into contact with the nuclear spins, which are then cooled. The rotation is repeated until the nuclear spin polarisation is steady.

30 The present invention can give the same information about the target compared to any previously known NMR method, but with the advantage of increased sensitivity. A further

advantage of this invention is that assay reagent containing an NMR active nucleus may in many cases, provide the same information previously provided by corresponding ^{14}C -labelled compounds, whilst being free from the problems associated with radioactive isotopes.

5 One further advantage according to the present invention is the increased signal-to-noise ratio. Another improvement with the present invention is that the time required to perform the assay is in general much shorter than the previously known methods. These improved parameters/results may be expressed as a "shortening effect", being the improvement of signal-to-noise ratio per unit time, and will be discussed further.

10 Yet another advantage compared e.g. with assays using fluorescent reagents is that there is no need to add an additional chemical component to the assay reagent to assist detection. There is always a disadvantage with techniques such as the fluorescent methods because the additional chemical component may influence the measurement.

15 The present invention provides an *in vitro* assay method which comprises:

- 20 a) using an assay reagent containing at least one NMR active nucleus to perform an assay, and
b) hyperpolarising at least one NMR active nucleus of the assay reagent;

Wherein steps a) and b) are performed simultaneously or sequentially in either order, and

- c) analysing the assay reagent and/or the assay by NMR, and
d) optionally using the NMR data obtained in step c) to generate further assay result(s).

25 As used herein, NMR active nuclei are those having non-zero nuclear spin and include ^1H , ^{13}C , ^{15}N , ^{19}F , ^{29}Si , ^{31}P and/or deuterium. Of these, ^{13}C and ^{15}N are preferred and ^{13}C is particularly preferred. Preferably the assay reagent for use in the assay according to this invention comprises an artificially-enriched abundance of an NMR active nucleus.

30 In a further preferred embodiment of the invention, the enriched compound comprises the artificially enriched NMR active nuclei, e.g. ^{13}C , at one specific position. Alternatively, in another preferred embodiment the compound comprises enriched NMR active

nuclei in 1-10 defined positions. A further alternative embodiment of the present invention is to have the assay reagent uniformly labelled with artificially enriched NMR active nuclei.

An assay reagent is a substance or compound that takes part in an assay, by being introduced as an initial reagent or by being formed *in situ* and perhaps transiently during the assay, or by being formed as a product of the assay. An assay is a test performed partly or wholly *in vitro* in which a physical or chemical change involving a biological species is observed. This change may have occurred both *in vivo* and *in vitro*. A biological species is one which is present in living systems or which is introduced into and is reactive with such systems. Preferred assay methods covered by this invention are related to biological macromolecules such as proteins (e.g. enzymes, receptors, DNA, and RNA binding proteins, carrier proteins), oligonucleotides (e.g. DNA and RNA probes, DNA and RNA consensus sequences), macrocyclic molecules (e.g. cyclodextrin) carbohydrate macromolecules and lipids.

Many assays involve a reaction in which a chemical bond is broken. According to another embodiment of the present invention, the assay reagent is an organic compound comprising one or more NMR active nuclei wherein these nuclei are associated with a bond which is broken during the course of the assay.

According to another embodiment of the present invention, the assay reagent contains two or more different types of NMR active nuclei, e.g. both ^{13}C and ^{15}N . Each active nucleus produces a distinct NMR spectrum and when the assay method is performed its results in changes to the chemical and/or physical environment of the nucleus. The changes to the environment are mirrored by spectral changes, which can be monitored.

The degree of hyperpolarisation of the NMR active nucleus covered by this invention is in excess of 0.1 %, more preferably 1 % and even more preferably at least 10 % above the equilibrium population of the excited state.

Surprisingly, assay methods where even smaller enhancement is achieved may effectively be performed due to the shorter time needed for the total assay measurement. One

important aspect of the present invention is thus an assay wherein the time required to give a defined signal-to-noise is considerably shortened by the use of this hyperpolarisation technique compared to known assay techniques without hyperpolarisation. The shortening effect is expressed as the improvement of signal-to-noise ratio per unit time, dB $\sqrt{\text{Hz}}$. This effect is preferably a factor of 10 or more, more preferably a factor of 25 or more and even more preferably a factor of 50 or more. In some embodiments, this effect is particularly a factor of 200 or more or even a factor of 1000 or more.

The assay can be carried out with the NMR active nucleus in the assay reagent already hyperpolarised. Alternatively, the assay may be carried out and the NMR active nucleus subsequently hyperpolarised prior, or at the same time, as the assay/assay reagent is analysed by NMR spectroscopy. Whilst the first arrangement enables real time studies of the assay to be carried out, this is often not necessary and, in these circumstances, the second method is very useful. As hyperpolarisation of the NMR active nucleus will sometimes be carried out at a low temperature, e.g. 20 K or less, the assay can be started and then effectively frozen by lowering the temperature. The assay/assay reagent is then hyperpolarised and analysed by NMR spectroscopy. By carrying out this process a number of times, either on the same assay or on parallel assays, a series of "snap-shots" of how the assay is proceeding may be obtained.

When hyperpolarisation is effected by exchange in solution phase, the hyperpolarising agent can be introduced as one batch, continuously or intermittently. Some conditions would lead to rapid disappearance of the hyperpolarisation. However, continuous or intermittent hyperpolarisation will give adequate signal intensity. Repeating the hyperpolarisation – acquisition sequence will also enhance the signal to noise ratio.

Agents, such as organic solvents, may in some situations be added to the assay, and/or to the NMR active nucleus if this is to be hyperpolarised prior to the assay, in order to prolong the life time of the hyperpolarised NMR active nucleus in the assay reagent, without interfering with the assay reagent and/or assay method.

Assays can be carried out by quantifying the appearance, or the continued presence, or the disappearance of spectral patterns. For example, on binding or hybridisation of an assay

reagent the chemical shift of the signals derived from the NMR active nucleus in the assay reagent will change. The different relaxation times of the different NMR active nuclei need to be taken into account if the quantification measurement is to be accurate.

5 It will be apparent to those skilled in the art that some NMR active nuclei, also referred to herein as hyperpolarisable atoms, retain their hyperpolarisation for a longer period than others at a given set of physical parameters. Thus, the order in which steps (a) and (b) of the method are carried out may, to some extent, be determined by the choice of NMR active nucleus. Whilst there may be advantages in carrying out the hyperpolarisation of the assay reagent and then
10 monitoring its NMR spectrum during the reaction, it is possible to "freeze" the reaction at any time. This may be achieved by reducing the temperature after the assay reagent has been added and then hyperpolarising the NMR active nucleus and comparing the spectra obtained with that of the assay reagent in a state where it has not undergone biological or chemical reaction(s).

15 As used herein, NMR active nuclei are those having non-zero nuclear spin and include ^1H , ^{13}C , ^{15}N , ^{19}F , ^{29}Si , ^{31}P and deuterium. Of these, ^{13}C and ^{15}N are preferred and ^{13}C is particularly preferred. ^{13}C is present at a natural abundance (relative to ^{12}C) of about 1%. Just as the labelling of organic compounds with radioactive ^{14}C is widely practised, so compounds, e.g. organic compounds can be labelled or enriched with ^{13}C , either generally or at specific positions
20 in the molecule. Preferably, the organic compounds for use in the assay according to this invention comprise an artificially-enriched abundance of ^{13}C , either generally or at least in one specific position, at an abundance of at least 5%, suitably at least 10%, more suitably at least 50%, preferably at least 75%, more preferably at least 90% and ideally at approaching 100%.

25 The present invention also covers the use of compounds comprising an artificially-enriched abundance of ^{15}N of at least 1%, suitably at least 5%, more suitably at least 10%, preferably at least 50% and more preferably at least 75% or more, and ideally at approaching 100%.

30 For assay reagents comprising ^{29}Si the preferred level of artificially enriched abundance is at least 10% and more preferred at a level of 50% or more, even more preferably at least 75 % or more and ideally at approaching 100 %.

For achieving as long T_1 as possible, the enriched compounds in some methods covered by the invention are preferably those in which the NMR active nucleus is surrounded by a double bond or one or more non-MR active nuclei such as O, S and/or C. In some cases, nearby
5 protons to the NMR active nucleus may be substituted by deuterium.

In one embodiment of the invention, step c) is performed by examining the assay reagent using both NMR spectroscopy to obtain spectral data from one or more discrete physical locations and repeating the examination at least once so as to obtain quantitative information
10 about kinetic or time-dependant alteration in chemistry, environment or structure of the assay reagent.

Assays envisaged according to this invention include for example, competition assays (e.g. receptor-ligand antagonism, enzyme-substrate inhibitors, protein-protein interaction
15 inhibitors), binding assays (e.g. receptor-ligand agonism, enzyme-substrate reactions, protein-protein interactions), immunoassays (e.g. for specific analytes), hybridisation assays (e.g. nuclease assays, mutation analysis, mRNA and DNA detection), tests involving cells, organs and/or whole organisms. Thus, the invention covers binding studies performed on tissue sections, cultured cells, cellular metabolites, micro-organisms and macro-organisms. Preferred examples
20 are discussed in the following paragraphs. Labelling with an NMR active nucleus where each molecule may be labelled at one or more chemical positions, will allow unique NMR assignments of e.g. starting material, intermediates and products of a biological reaction. Thus dual, triple etc labelling experiments can be carried out and 'stop-flow' measurements made with identical chemical species. For example, theoretically, all the six carbon atoms in glucose could be
25 individually or collectively replaced by ^{13}C , so that one to six of the carbon atoms are ^{13}C which can be hyperpolarised. Each hyperpolarised ^{13}C will give rise to a chemical shift, which will be specific to that individual carbon and different to other ^{13}C positions in the molecule, i.e. C-1 will be different from C-2, etc.

A non-exclusive list of the types of molecules into which NMR active nuclei may be incorporated includes:

(i) Amino acids

Amino acids contain carbon, nitrogen and hydrogen and therefore any amino acid can be labelled at a single or multiple positions with one or more different NMR-active nuclei.

(ii) Lipophilic compounds

These would contain fatty acids, phospholipids, glycerol, cholesterol and its esters, sphingosine and its esters. These contain carbon, hydrogen and in some cases nitrogen and/or phosphorus and can be labelled at a single or multiple positions with one or more different NMR-active nuclei.

(iii) Vitamins

These include the water-soluble and fat-soluble categories of essential nutrients. These contain carbon, hydrogen and in some cases nitrogen and/or phosphorus and can be labelled at a single or multiple positions with one or more different NMR active nuclei.

(iv) Nucleic acids etc

DNA contains the bases adenine, cytosine, guanine and thymine and their nucleosides and nucleotides. RNA contains the bases adenine, guanine, cytosine and uracil and their nucleosides and nucleotides. These contain carbon, hydrogen, nitrogen and in some cases phosphorus and can be labelled at a single or multiple positions with one or more different NMR-active nuclei.

In one preferred embodiment of the invention the hyperpolarisation transfer is achieved by using a hyperpolarised noble gas, or a mixture of such gases, to effect nuclear polarisation of an assay reagent comprising at least one NMR active nucleus other than the noble gas.

5

When the hyperpolarisation of the assay reagent is achieved by an artificially enriched hyperpolarised noble gas, the hyperpolarised noble gas is preferably ^3He or ^{129}Xe . Such isotopically enriched gases are now commercially available at high isotope purity and can be polarised to a high degree of hyperpolarisation. The hyperpolarised gas may, if desired, be stored
10 for extended periods of time in the polarised state, by keeping the gas at very low temperatures, especially in a frozen form.

A hyperpolarised noble gas may be used in step b) of the present invention to effect nuclear polarisation of an assay reagent comprising at least one NMR active nucleus other than
15 the noble gas. The hyperpolarised gas may be in the gas phase, condensed or may alternatively be liquid e.g. by being dissolved or emulsified in a lipophilic solvent such as a lipid or a fluorocarbon solvent, or in a suspension or a solid e.g. by being adsorbed or frozen on to a solid surface. In some cases, liposomes or microbubbles may encapsulate the hyperpolarised noble gas.

The assay reagent may be solid, semi-solid or fluid. A hyperpolarised gas may be
20 bubbled into a fluid assay system. Alternatively, a hyperpolarised gas solution may be mixed with a fluid assay. The hyperpolarised gas may be cooled and/or maintained in a magnetic field to preserve the hyperpolarisation. Similarly the resulting assay reagent comprising at least one polarised NMR active nucleus may preferably be cooled and/or maintained in a magnetic field in
25 order to preserve the polarisation and/or facilitate polarisation transfer.

One advantage with hyperpolarisation transfer by ^3He or ^{129}Xe is that these gases are essentially chemically inert and will not adversely affect the assay reagent or the assay. In addition, as in gaseous form, ^3He and/or ^{129}Xe are easily separated from the assay medium,
30 permitting facile repeat studies.

In one embodiment, a flow of hyperpolarised gas in the liquid state at elevated pressure and/or low temperature is passed through a column of the assay reagent. The gas may be pumped off and the process repeated until a suitable level of polarisation is achieved.

Alternatively, a hyperpolarised gas is frozen/crystallised on the solid/frozen surface of the solid assay reagent. This compound may preferably have been prepared with as large a surface area as possible, e.g. as a finely divided powder.

In some cases, it is desirable to remove part of or substantially the whole of the hyperpolarisable gas from the assay reagent/system as rapidly as possible. If desired, the gas may be reused which may be important due to the expense of isotopically-enriched noble gases. Many physical and chemical separation or extraction techniques known in the art may be employed to effect rapid and efficient separation of the hyperpolarised gas and the assay system.

In one embodiment of the invention when performed with the assay reagent in the solid phase, it is especially important that the content of ^{131}Xe should be as low as possible. The preferred content of ^{131}Xe is thus below 0.5 % of the total Xe content, and more preferably below 0.05 %.

In a further aspect, the present invention provides a method for optimising the polarisation enhancement factor when the assay reagent is hyperpolarised by a noble gas in solution. Thus the enhancement of the target nuclear spin can be optimised by slowing the dynamics of the molecules (atoms) in the solution. The dynamics can be slowed down, e.g. by increasing the viscosity of the solvent. The polarisation enhancement factor may also depend on the concentration of the noble gas in the solution and the enhancement factor may be optimised further by adjusting the pressure and temperature.

The relaxation mechanisms and also the relaxation of the target nucleus are partly functions of the viscosity of the solvent. For a specific system of interest we may choose the optimal viscosity of the medium that will lead to the maximal polarisation enhancement factor of the target nucleus. The viscosity is determined by choice of solvent and temperature. Preferably, the viscosity should be at least 1000 mPs, more preferably at least 10000 mPs and especially preferably at least 100000 mPs.

In one embodiment of the invention, when the polarisation transfer occurs in solution, the pressure of xenon is as high as possible, preferably higher than $5 \times 10^5 \text{ N/m}^2$ (5 bar), more preferably higher than $5 \times 10^6 \text{ N/m}^2$ (50 bar), even more preferably higher than $1 \times 10^7 \text{ N/m}^2$ (100 bar) and particularly higher than $2 \times 10^7 \text{ N/m}^2$ (200 bar). However, the pressure must never be so high so that the biological molecule will be totally or partly adversely effected.

It is preferred that the solvent comprises as few atoms which possess magnetic moment as possible and is as low magnetogyric ratio as possible. The transfer of polarisation in a highly viscous medium may be followed by solution spectroscopy under high-viscosity conditions (broad lines).

Alternatively, the viscosity may be lowered prior to spectroscopy, either by a change in temperature or by a change in the chemical composition of the solvent. If the high-viscosity medium is formed by a pH-sensitive gel-forming agent, then the viscosity might be lowered e.g. by a change in pH. Changes of temperature, ion-strength as well as the use of specific additives may also be considered.

In a further embodiment, the present invention provides a method wherein the hyperpolarisation transfer is effected by use of a very high field and with very low temperature (Brute force). The magnetic field strength used should be as high as possible, suitably higher than 1T, preferably higher than 5T, more preferably 15T or more and especially preferably 20T or more. The temperature should be very low e.g. 4.2K or less, preferably 1.5K or less, more preferably 1.0K or less, especially preferably 100 mK or less.

25

US 5479925 discloses a method for generating MR angiograms in which a contrast agent is passed through a small, high field polarising magnet *in vitro* in order to generate a high longitudinal magnetisation in the agent prior to its administration to the subject. However, there is no mention of the use of an enriched NMR active nucleus. When this Brute force method is used, and thermodynamic equilibrium is attained, all nuclei in the assay reagent will be highly polarised relative to room temperature and to normal magnetic fields used in MRI.

30

A major practical problem when using this technique is the time required for the thermal equilibrium to occur. However, the Brute force embodiment may be modified in order to solve this problem as described below.

5 It is possible to use a technique of low-field matching to increase the relaxation rate and the degree of polarisation of the nuclear spins in solids at low temperature. This has the additional advantage that equipment used in the Brute force polariser does not need to possess any radio frequency electronics.

10 A way of speeding up the polarisation of the NMR active nuclei, at least for ^{13}C and ^{15}N and at the same time obtaining a better polarisation is to use cross-polarisation from the quickly relaxing proton to the slowly relaxing carbon, a method routinely used in solid-state NMR spectroscopy. The situation may be further improved by utilising the procedure of spin locking under Hartman-Hahn conditions. However, radiofrequency electronics are required and
15 furthermore the homogeneity of the magnetic field must be high enough to allow precise pulse angles. A simplified method to allow for thermal contact between the protons and the NMR active nucleus (e.g. ^{13}C or ^{15}N) is to remove the assay from the magnet for a fraction of a second and repeat this procedure after the protons have repolarised, successively building up the polarisation until the spin-temperature of the two nuclei become the same.
20

A further improvement of the Brute force embodiment of this invention is to optionally expose the assay system to a relaxation shortening effect in order to attain thermodynamic equilibrium at said low temperature. The relaxation shortening effect may be provided by exposure to field cycling to a field allowing cross polarisation, gradually increasing
25 the magnetic field at such a rate that the increase in polarisation of the assay reagent is maximised. This effect may also be achieved by adding magnetic material to the assay reagent during the period when the assay reagent is exposed to low temperature.

In a further embodiment, the present invention provides a method for the polarisation
30 transfer using the DNP method effected by a DNP agent, to effect nuclear polarisation of an assay reagent comprising at least one NMR active nucleus. In the solid phase, there are two aspects of DNP, namely "the solid effect" and the thermal mixing.

Most known paramagnetic compounds may be used as a "DNP agent" in this embodiment of the invention, e.g. transition metals such as chromium ions or organic free radicals such as nitroxide radicals and trityl radicals (WO 98/58272). Where the DNP agent is a paramagnetic free radical, the radical may be conveniently prepared *in situ* from a stable radical precursor by a radical-generating step shortly before the polarisation, or alternatively by the use of ionising radiation. Energy, normally in the form of microwave radiation, is provided in the process which will initially excite the paramagnetic species. Upon decay to the ground state, there is a transfer of polarisation to an NMR active nucleus of the target material. The method may be conveniently carried out by using a first magnet for providing the polarising magnetic field and a second magnet for providing the primary field for MR spectroscopy/imaging.

In some cases, the radical will be non-reusable and may conveniently be discarded after use. Many physical and chemical separation or extraction techniques are known in the art, which may be used if it is desirable to remove the DNP agent from the assay system in a rapid and/or efficient separation step. Magnetic properties may e.g. be used to achieve the separation. It is particularly preferred to use a heterogeneous system, e.g. a two-phase liquid, a solid in liquid suspension or a high surface area solid substrate within a liquid. For any heterogeneous system, separation may be achieved by e.g. filtration, decanting, chromatographic or centrifugal methods.

In a further embodiment, the present invention provides a method wherein the polarisation transfer is achieved by exposing the assay reagent to para hydrogen-enriched hydrogen gas in the presence of a suitable catalyst. The assay reagents suitable for use are prepared from precursors which are able to be hydrogenated and which will therefore typically possess one or more unsaturated bonds, e.g. double or triple carbon-carbon bonds.

Hydrogen molecules exist in two different forms, para hydrogen ($p\text{-H}_2$) where the nuclear spins are anti parallel and out of phase (singlet state) and ortho hydrogen ($o\text{-H}_2$) where the spins are parallel or anti parallel and in phase (triplet state). At room temperature, the two forms exist in equilibrium with a 1:3 ratio of para:ortho hydrogen. However, preparation of para hydrogen enriched hydrogen can be carried out at low temperature, 160K or less, in the presence of a catalyst. The para hydrogen formed may be stored for long periods, preferably at low

temperature, e.g. 18-20K. Alternatively it may be stored in pressurized gas form in containers which have an inner surface which is non-magnetic and non-paramagnetic.

When the p-H₂ molecule is transferred to the precursors of the assay reagent (by
5 means of catalytic hydrogenation with e.g. (PPh₃)₃RhCl), the proton spins remain anti parallel
and begin to relax to thermal equilibrium with the normal constant T₁ of the hydrogen in the
assay molecule. However, during relaxation some of the polarisation may be transferred to
neighbouring nuclei by pulse sequence (Progress in Nuclear Spectroscopy, 31, (1997), 293-315),
low field cycling or other types of coupling. The presence of the NMR active nucleus as e.g. ¹³C
10 (and ¹⁵N etc) with a suitable substitution pattern close to the relaxing hydrogen may lead to the
polarisation being trapped in the slowly relaxing ¹³C (or ¹⁵N etc) resulting in a high enhancement
factor.

A further hyperpolarisation transfer embodiment of this invention is the spin
15 refrigeration method. This method covers spin polarisation of a solid assay by spin refrigeration
polarisation. The assay is doped with or intimately mixed with a suitable paramagnetic material
such as Ni²⁺, lanthanide and actinide ions in crystal form with a symmetry axis of order three or
more. The instrumentation is simpler than that required for DNP with no need for a uniform
magnetic field. The process is carried out by physically rotating the sample around an axis
20 perpendicular to the direction of the magnetic field. The prerequisite for this to work is that the
paramagnetic species has a highly anisotropic g-factor.

Hybridisation assays are very widely used for sequencing and for detection of point
or deletion mutations in nucleic acids. When a conventionally labelled polynucleotide probe is
25 hybridised with a polynucleotide target, analysis of the melting temperature or other property of
the hybrid can give some limited information about the nucleotide sequence of the target.

The present invention can give the same information about the target compared to
any previously known NMR methods available, but with the advantage of increased sensitivity. A
30 polarised NMR active nucleus generates an NMR spectrum which is dependent on its
environment, i.e. the atoms surrounding the NMR active nucleus, both intramolecular (atoms
within the same molecules as the NMR active nucleus) and intermolecular (atoms in the other

molecules nearby the NMR active nucleus). The environment thus extends beyond the labelled molecule itself to other molecules in the immediate vicinity. Thus for example, a nucleotide labelled with polarised NMR active nucleus, e.g. ^{13}C and/or ^{15}N , when incorporated into a single stranded polynucleotide chain, can give information about two or more adjacent nucleotide
5 residues in the chain. When that labelled polynucleotide probe is hybridised with a polynucleotide target, NMR spectroscopic analysis of the NMR ^{13}C label can give information about the complementary nucleotide residue in the target.

In one embodiment of the present invention, comparative and/or parallel testing is
10 performed to maximise the information available from the NMR measurements.

Biological macromolecules such as nucleosides or nucleotides or nucleotide analogues can readily be enriched with a NMR active nucleus, e.g. ^{13}C and/or ^{15}N at one or several specified points in the molecule. Polarisation of the NMR active nucleus, e.g. ^{13}C ,
15 preferably by contact with a hyperpolarised noble gas, may be effected either before, during or after incorporation of the monomer into a polynucleotide; and before, during or after hybridisation of that polynucleotide with a complementary strand.

Figure 1 demonstrates a hybridisation assay in which the use of an oligonucleotide or
20 polynucleotide is used to detect the presence of single nucleotide polymorphisms (SNPs) in a gene, or fragment of a gene. An oligonucleotide or polynucleotide probe is prepared in which one or more of the atoms has been replaced by a hyperpolarisable isotope, e.g. ^{13}C , ^{15}N or ^1H . This probe is then hybridised to the gene or the gene fragment. The probe will be "targeted" to information-rich parts of the gene and may be selected so that the probe binds only to that part of
25 the DNA containing a specific mutation, or, potentially, more than one mutation. If desired, a set of probes, each probe containing a hyperpolarisable isotope, can be added to a gene or gene fragment, each probe being targeted to a different part of the gene/gene fragment. As each probe will have a characteristic chemical shift by NMR spectroscopy, the spectrum of the mixture of the probes with the target can be taken and resolved to indicate which probes have bound and
30 which have not.

The probe may be polarised before, during, or after hybridisation and a determination

carried out by NMR of whether a shift has occurred in the signal obtained from the hyperpolarised isotopic atom(s). If a shift has occurred, then the probe is (by inference) in a different chemical environment indicating hybridisation. Clearly information can be obtained from both positive and negative results, e.g. a probe could be constructed from the “natural” gene, a naturally occurring DNA sequence, and if results indicate that this has failed to bind, probes could be tested containing anticipated mutations. This technique facilitates itself to use of an array-type format in which a number of hyperpolarisable probes are used in the assay which each vary by one nucleotide. The identity of the SNP can be determined by the hybridisation pattern of the probes to the gene/gene fragment.

As mentioned earlier, many assays involve a reaction in which a chemical bond is broken. According to one embodiment of the present invention, the assay reagent is an organic compound comprising one or more NMR active nuclei associated with a bond which is broken during the course of the assay. In the case of a single NMR active nucleus, this is located preferably at the actual site of the breaking of the chemical bond such that the change in local environment of the active nucleus subsequent to the bond breaking will give rise to a significant change in the spectrum of the NMR active nucleus. The NMR spectra of two or more active nuclei will be different, depending on whether they are present within the same molecule or in different molecules. When two or more NMR active nuclei are in an appropriate proximity to one another they are said to be spin coupled. This gives rise to a distinct NMR spectrum which can be monitored. It is therefore possible to analyse by NMR spectroscopy the rate and extent of the bond breaking by the disruption of the spin coupling. In this and other assays, the assay reagent may be analysed repeatedly by NMR spectroscopy at known time intervals so as to generate information about a change over time of the assay reagent.

Figure 2 demonstrates a proteolysis assay. The starting substrate for the reaction contains two hyperpolarisable isotopes, in this case ^{13}C , which are sufficiently close together, either by virtue of being reasonably adjacent in the chain of amino acids comprising the molecule, or by the 3-dimensional conformation of the molecule held in a “conformational lock”. In these situations, NMR spectra J coupling (scalar coupling) of the signal occurs and the NMR spectra of the molecule is recorded.

The molecule is then brought into contact with an enzyme capable of altering the chemical composition of the substrate. If cleavage occurs between the amino acids containing the hyperpolarisable isotopic atoms, then the J coupling and the chemical shift values change which will be observed by NMR spectroscopy and/or NMR imaging. Two new spectra will appear, one
5 for each of the individual cleavage products. If there is no cleavage, the original spectrum remains.

A similar assay can be carried out where the starting substrate is a chain of nucleotides and the cleavage enzyme an endonuclease.

10 In another aspect of the invention, an assay reagent may be administered to a macro-organism, e.g. a human or animal, and NMR spectroscopic analysis performed of blood, excreta, e.g. urine, faeces or breath, or samples of the macro-organism.

15 In yet another aspect of the invention, an assay reagent may be used in binding studies on bacteria or other eukaryotic or prokaryotic micro-organisms or cultured cells.

Assays according to one embodiment of this invention may conveniently be carried out in multiwell plates. An assay reagent in each well may e.g. be hyperpolarised by contact with a
20 hyperpolarised noble gas, prior to addition of other assay reagents. Alternatively, an assay reagent in bulk may be hyperpolarised with a hyperpolarised noble gas prior to being dispensed into individual wells of a multiwell plate. In many cases, assays can be performed in a homogenous mode, that is to say without the need for a separation step to remove one fraction of the labelled reagent.

25 In addition, in cases where the spectra of the ^{13}C labelled assay components are distinct from one another, more than one assay may be performed and simultaneously monitored in a single well or spot of a multi-assay array. This would allow multiplexing of several related or unrelated assays in parallel within a single well or spot in a multi-assay array which is either
30 ordered or random. In addition the technique may be applied to aerosol droplets where no well, container or surface is used to contain the assay and to analysis of samples in flow-through devices.

Figure 3 illustrates how the incorporation of a material (for example an amino acid) into a cell can be measured. The material incorporates a hyperpolarisable isotopic atom, in this case ^{13}C . Its NMR spectrum in a hyperpolarised state in the media used for the experiment is recorded. If the material crosses the cell membrane then the environment in which the material finds itself will change and this will affect the NMR chemical shift of the material. The precise chemical shift will depend on the environment of the material within the cell, for example it may be possible to identify whether it has crossed into the cell nucleus. Alternatively, the material may be bound to the surface of the cell, again a different spectrum will result. In addition, metabolites that contain a hyperpolarisable isotopic atom may be detected either inside the cell or after they are excreted from this. The spectra obtainable on these metabolites can be used for their identification and/or to give information on their structure.

In one embodiment of the present invention, the assay is performed at a relatively cold temperature. However, in some situations the assay is carried out at room temperature.

It is important that the probes, vials, coils etc are coated or made of materials which do not induce loss of polarisation, such as para-magnetic nuclei. Preferred such materials are e.g. plastic, aluminium, Teflon and glass (with low iron content) materials. A further embodiment of the method according to the invention is thus the use of materials such as aluminium, plastic, glass and/or Teflon for the wells, vials, containers and any coils. A metal may also be used coated with a non-para magnetic oxide layers (e.g. Ti, Mg or Ag).

In one preferred embodiment of the invention, the assay is carried out in an NMR tube, with a gas-tight seal, permitting the addition (and/or removal) of a hyperpolarised gas to (and/or from) the assay reagent.

A variety of NMR spectroscopy and/or NMR imaging manipulation methods may be used, e.g. magic angle spinning and pulse sequence like WAHUHA or MLEV-8 to obtain high resolution spectrum when the assay reagent is a solid or semi-solid state.

A further embodiment of the present invention is an *in vitro* kit for carrying out the assay method as defined. The kit comprises a well, vial or any other suitable container comprising one or more assay reagents optionally together with additives wherein the hyperpolarisation transfer occurs. One embodiment of the invention concerns an *in vitro* kit where the NMR analysis of step (c) of claim 1 is carried out in the same well, vial or container as the polarisation transfer is carried out.

The invention is illustrated with reference to the following non-limiting example. Modifications of the method according to this example include the addition of the noble gas directly into the spectrometer and the use of different pulse techniques.

Example 1.

Polarisation transfer from hyperpolarised ^{129}Xe to the singly labelled peptide AcYRARV(F, ^{13}C -amide)FVRAAK-NH₂

Hyperpolarized ^{129}Xe was generated by optical pumping as described by B.Driehuys et al., Appl.Phys.Lett. 69 (12), 1996. The isotopic composition of the gas was 80% ^{129}Xe and 0.25% ^{131}Xe (the rest non-magnetic isotopes of Xe). The degree of polarization was estimated to be $10\% \pm 3$.

The freeze-dried peptide (3.4 mg) was placed in an ordinary 5 mm thin-walled NMR-tube. The glass tube was connected to the outlet of the polarizer by means of 60 cm of plastic tubing. The tube was evacuated and then filled with nitrogen four times.

The hyperpolarized gas was generated and collected on a cold finger at liquid nitrogen temperature in a holding field of 200 mT over a period of 15 minutes which is estimated to give a volume of 50 ml of Xenon at NTP. A narrow Dewar vessel with liquid nitrogen was placed in a magnet with a field strength of 0.3 T. The collected xenon was thawed and gradually refrozen on the peptide from the bottom and up by gradually lowering the tube into the liquid nitrogen bath. The system was then filled with helium to one atmosphere. The sample, with the plastic tubing still connected but open to the surroundings, in the Dewar in the 0.3 T magnet with the poles in horizontal configuration was then moved into the stray-field of the 7 T magnet (vertical polarity)

of an NMR-spectrometer. The sample was then rapidly transferred to the spectrometer and was in the process subjected to a minimum magnetic field of 0.3 mT.

A ^{13}C spectrum was recorded with a spectral window of 100 kHz and a broad ^{13}C signal
5 was obtained. The sample was then left to polarize in the magnet and a background signal was recorded overnight, and care was taken to allow for full relaxation between the pulses.

The enhancement was measured to 6 ± 1 times the thermodynamic equilibrium at 7 T and 291 K.

10

The time from the beginning of freezing the xenon in the NMR tube to the acquisition of the spectrum was 5 minutes.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2

C L A I M S

1. An *in vitro* assay method which comprises:
 - a) using an assay reagent containing at least one NMR active
 - 5 nucleus to perform an assay, and
 - b) hyperpolarising at least one NMR active nucleus of the assay reagent;
wherein steps (a) and (b) are performed simultaneously or sequentially in either order, and
 - c) analysing the assay reagent and/or the assay by NMR and
 - 10 d) optionally using the NMR data obtained in step c) to generate further assay result(s).
2. The method of claim 1 wherein the NMR active nucleus is ^{15}N , ^{19}F , ^{31}P , ^1H , ^{29}Si and/or ^{13}C .
3. The method of claim 2, wherein the NMR active nucleus is ^{15}N or ^{13}C .
4. The method of any of claims 1 to 3, wherein the assay reagent is a compound which contains an artificially high concentration of an NMR active nucleus.
5. The method of claim 4, wherein the assay reagent contains an artificially high concentration in 1-10 defined positions.
6. The method of any of claims 1 to 5, wherein the assay reagent is an organic compound comprising one or more NMR active nuclei associated with a bond which is broken during the course of the assay.
7. The method of claim 6, wherein the assay reagent contains two or more NMR active nuclei and each NMR active nucleus produces a distinct NMR spectrum and when the


assay method is performed, it results in changes to the chemical and/or physical environment of the nucleus and this is mirrored by spectral changes which can be monitored.

- 5 8. The method of any claim 1-7, wherein the assay reagent is analysed repeatedly in step c) at known time intervals so as to generate information about a change with time of the assay reagent.
9. The method of any one claim 1 to 8, wherein the assay reagent is a Nucleotide, or
10 nucleotide analogue, polynucleotide, amino acid analogue, polypeptide or protein.
10. The method of any one of claims 1 to 9, wherein the assay is a nucleic acid hybridisation assay.
- 15 11. The method of any one of claims 1 to 10, wherein the assay is a binding assay.
12. The method of claims 1 to 11, wherein the assay reagent is a compound specifically labelled with at least one NMR active nucleus and the assay reagent is administered to a micro-organism, macro-organism or cultured cells, cellular metabolites or an excretion
20 product of the assay reagent are hyperpolarised and analysed by nuclear magnetic resonance spectroscopy, nuclear magnetic resonance imaging or both.
13. The method of claims 1 to 12, wherein the assay is a binding study performed using micro-organisms or cultured cells.
- 25 14. The method of claims 1 to 13 wherein the hyperpolarisation transfer is repeated to enhance the signal-to-noise ratio.
- 30 15. The method of claim 1 to 14 wherein the shortening effect as expressed by the improvement of signal-to-noise per unit time is a factor of 10 or more compared to known assay techniques without hyperpolarisation.

16. The method of claims 1 to 15 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer from a hyperpolarised noble gas, or a mixture of hyperpolarised noble gases.
- 5 17. The method of claim 16 wherein the noble gas is ^{129}Xe .
18. The method of claim 16 wherein the noble gas is ^3He .
19. The method of claims 16 to 18 wherein the hyperpolarisation is transferred by a
10 hyperpolarised noble gas in solution and wherein the viscosity of the solution is at least 1000 mPs.
20. The method of claims 1 to 15 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer at a temperature of 4.2 K or less in
15 the presence of a magnetic field of at least 1 T.
21. The method of claims 1 to 15 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by polarisation transfer using dynamic nuclear
20 polarisation.
22. The method of claims 1 to 15 where the hyperpolarisation of the NMR active nucleus of the assay reagent is carried out by para hydrogen induced polarisation.
23. The method of claims 1 to 15 where the hyperpolarisation of the NMR active nucleus of
25 the assay reagent is carried out with the spin refrigeration technique.
24. The method of claims 1 to 23, wherein more than one assay is multiplexed and monitored by NMR spectroscopy and/or NMR imaging.
- 30 25. The method of claims 1 to 24 wherein the assay is performed in a multiwell or multispot assay array.

26. The method of claims 1 to 25 wherein step c) is performed by examining the assay reagent using both NMR spectroscopy to obtain more than one spectrum, and magnetic resonance imaging to obtain one or more discrete spectral location, and repeating the examination at least once so as to obtain quantitative information about kinetic or time-
5 dependant alteration in chemistry, environment or structure of the assay reagent.
27. The method of claim 1 to 26, wherein step c) is performed in an aerosol or flow-through device applied to aerosol droplets where the well, surface or container is used to contain the assay reagent.
- 10 28. An *in vitro* assay kit for carrying out the assay method as defined in claim 1 to 27 which comprises: one or more assay reagents each containing at least one NMR active nucleus contained in a well or vial or other suitable container for carrying out the hyperpolarisation of step (b) of claim 1.
- 15 29. The *in vitro* kit of claim 28 where the NMR analysis of step (c) of claim 1 is carried out in the same well, vial or container as the hyperpolarisation transfer is carried out.

1/2



2/2

Fig.2.



Spin dipole spectra2 Individual spectra

Fig.3.



Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)
Approved for use through 9/30/00 OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

**DECLARATION FOR UTILITY OR
DESIGN
PATENT APPLICATION
(37 CFR 1.63)**

☐ Declaration Submitted with Initial Filing
OR
☒ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

Attorney Docket Number PA-9848

First Named Inventor Knox

COMPLETE IF KNOWN

Application Number 09 / 869,629

Filing Date 28-Jun-2001

Group Art Unit To be assigned

Examiner Name To be assigned

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

NMR Spectroscopic In Vitro Assay Using Hyperpolarization

the specification of which (Title of the Invention)

☐ is attached hereto
OR

☒ was filed on (MM/DD/YYYY) 06/28/2001 as United States Application Number or PCT International

Application Number 09/869,629 and was amended on (MM/DD/YYYY) (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
9828852.5	Great Britain	12/30/1998	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
9918096.0	Great Britain	08/02/1999	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231

Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

+

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet
Page 1 of 2

Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
2-0 Neil				Cook			
Inventor's Signature				Date			
Residence: City		State		Country		GB	
Post Office Address		Tutshill Lodge, Beachley Road, Tutshill					
Post Office Address		Chepstow, Great Britain NP6 7EG GBN					
City		State		ZIP		Country	
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
3-0 Klaes				Golman			
Inventor's Signature				Date			
Residence: City		State		Country		SE	
Post Office Address		Nykcomed Innovation AB, Ideon Malmo					
Post Office Address		Per Albin Hanssons vag 41, S-205 12 Malmo Sweden SE					
City		State		ZIP		Country	
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
4-0 Haukur				Johannesson			
Inventor's Signature		Haukur Johannesson		Date		August 27, 2001	
Residence: City		State		Country		SE	
Post Office Address		Nykcomed Innovation AB, Ideon Malmo					
Post Office Address		Per Albin Hanssons vag 41, S-205 12 Malmo Sweden SE					
City		State		ZIP		Country	

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231 DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO Assistant Commissioner for Patents, Washington, DC 20231

+



Please type a plus sign (+) inside this box → +

PTO/SB/02A (3-97)
Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet
Page 1 of 2

Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Neil				Cook			
Inventor's Signature				20/8/01 Date			
Residence: City	Princeton	State	NJ	Country	USA	Citizenship	GB
Post Office Address	Amersham Pharmacia Biotech Inc, 800 Centennial Avenue						
Post Office Address	Piscataway, NJ 08855-1327, USA						
City		State		ZIP		Country	
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Klaes				Golman			
Inventor's Signature				Date			
Residence: City		State		Country	SE	Citizenship	DK
Post Office Address	Nycomed Innovation AB, Ideon Malmo						
Post Office Address	Per Albin Hanssons vag 41, S-205 12 Malmo Sweden						
City		State		ZIP		Country	
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])				Family Name or Surname			
Oksar				Axelsson			
Inventor's Signature				Date			
Residence: City		State		Country	SE	Citizenship	SE
Post Office Address	Nycomed Innovation AB, Ideon Malmo						
Post Office Address	Per Albin Hanssons vag 41, S-205 12 Malmo Sweden						
City		State		ZIP		Country	

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.



Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)
Approved for use through 9/30/98 OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet
Page 1 of 2

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Neil		Cook			
Inventor's Signature				Date	
Residence: City	State	Country	GB	Citizenship	GB
Post Office Address	Tutshill Lodge, Beachley Road, Tutshill				
Post Office Address	Chepstow, Great Britain NP6 7EG				
City	State	ZIP	Country		
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Klaes		Golman			
Inventor's Signature				Date	8-8-2001
Residence: City	State	Country	SE	Citizenship	DK
Post Office Address	Nycomed Innovation AB, Ideon Malmo				
Post Office Address	Per Albin Hanssons vag 41, S-205 12 Malmo Sweden				
City	State	ZIP	Country		
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Oksar		Axelsson			
Inventor's Signature				Date	3/8 2001
Residence: City	State	Country	SE	Citizenship	SE
Post Office Address	Nycomed Innovation AB, Ideon Malmo				
Post Office Address	Per Albin Hanssons vag 41, S-205 12 Malmo Sweden				
City	State	ZIP	Country		

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231


Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)
Approved for use through 9/30/98 OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet
Page 2 of 2

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Jan Henrik		Ardenkjaer-Larsen			
Inventor's Signature				Date	27-7-2001
Residence: City		State		Country	SE
				Citizenship	DK
Post Office Address	Nycomed Innovation AB, Ideon Malmo				
Post Office Address	Per Albin Hanssons vag 41, S-205 12 Malmo Sweden				
City		State		ZIP	
				Country	
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Inventor's Signature				Date	
Residence: City		State		Country	
				Citizenship	
Post Office Address					
Post Office Address					
City		State		ZIP	
				Country	
Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name (first and middle [if any])		Family Name or Surname			
Inventor's Signature				Date	
Residence: City		State		Country	
				Citizenship	
Post Office Address					
Post Office Address					
City		State		ZIP	
				Country	

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)
Approved for use through 9/30/00 OMB 0651-0032
Patent and Trademark Office, U. S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)
PCT/GB99/04410	12/23/1999	

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02A attached hereto

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact with the Patent and Trademark Office connected therewith: ☒ Customer Number 22840 OR ☐ Registered practitioner(s) name/registration number listed below

Name	Registration Number	Name	Registration Number

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto

Direct all correspondence to: ☒ Customer Number 22840 OR ☐ Correspondence address below

Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: ☐ A petition has been filed for this unsigned inventor

Given Name (first and middle if any)	Family Name or Surname
Peter	Knox

Inventor's Signature	<i>Peter Knox</i>	Date	30.7.01		
Residence: City	State	Country	GB	Citizenship	GB
Post Office Address	"Choppings", 34 Kings Road, Buckinghamshire				
Post Office Address	Chalfont St. Giles, Great Britain HP8 4HS				
City	State	ZIP		Country	

☒ Additional inventors are being named on the 2 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto